

Comments On The Sign of $\mathbf{CP}$ Asymmetries

I. I. Bigi ^a, A. I. Sanda ^b

^a*Physics Dept., Univ. of Notre Dame du Lac, Notre Dame, IN 46556, U.S.A.*

^b*Physics Dept., Nagoya University, Nagoya 464-01, Japan*

e-mail addresses:

bigi@undhep.hep.nd.edu, sanda@eken.phys.nagoya-u.ac.jp

Abstract

Several experiments are about to measure the $\mathbf{CP}$ asymmetry for $B \rightarrow \psi K_S$ and other decays. The standard model together with the Kobayashi-Maskawa ansatz for $\mathbf{CP}$ violation predicts the sign as well as the magnitude for this asymmetry. In this note we elucidate the physics and conventions which lead to the prediction for the sign of the asymmetry.

I. THE PROBLEM

It has been pointed out some time ago [1] that within the KM ansatz large **CP** asymmetries have to arise in B decays involving $B^0 - \bar{B}^0$ oscillations. In particular the channel $B_d \rightarrow \psi K_S$ combines a striking experimental signature with a clean theoretical interpretation [2]; therefore it is often referred to as the "golden mode".

A first experimental information on the **CP** asymmetry in it has recently become available [3,4]. The asymmetry is defined as follows

$$A_{\psi K_S} = \frac{\Gamma(\bar{B}_d(t) \rightarrow \psi K_S) - \Gamma(B_d(t) \rightarrow \psi K_S)}{\Gamma(\bar{B}_d(t) \rightarrow \psi K_S) + \Gamma(B_d(t) \rightarrow \psi K_S)} = \text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right) \sin(\Delta M_{B_d} t) \quad (1.1)$$

where $\bar{\rho}(\psi K_S)$ denotes the ratio of transition amplitudes

$$\bar{\rho}(\psi K_S) = \frac{A(\bar{B} \rightarrow \psi K_S)}{A(B \rightarrow \psi K_S)}. \quad (1.2)$$

while q and p relate the mass eigenstates to the flavour eigenstates B_d and $\bar{B}_d$ (see below).

Two groups found

$$\text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right) = \begin{cases} +(3.2^{+1.8}_{-2.0} \pm 0.5) & \text{OPAL Collaboration [3]} \\ +(1.8 \pm 1.1 \pm 0.3) & \text{CDF Collaboration [4]} \end{cases} \quad (1.3)$$

In reality the observable $\text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right)$ is bounded by -1 and +1. It falls outside this range in the data since they require subtracting a background of uncertain size. Thus the numbers have to be taken with a grain of salt. Even so they indicate that values close to +1 are very strongly disfavoured.

This raises the following question: Can one *predict* also the *sign* in addition to the size of the asymmetry?

At first one might think that to be impossible. For $A_{\psi K_S}$ is a product of $\text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right)$ and $\sin(\Delta M_{B_d} t)$, see Eq.(1.1). The observable sign of $A_{\psi K_S}$ thus depends on the signs of both $\text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right)$ and ΔM_{B_d} , and the sign of the latter cannot be defined nor determined *experimentally* in a feasible way – in contrast to the case with kaons.

In this note we will show that the overall sign of the asymmetry $A_{\psi K_S}$ – and likewise for other asymmetries – can be predicted within a given theory for $\Delta B = 2$ dynamics. For

those who have thought about this question carefully, this has been known for a long time. Indeed it has been discussed in [5]. Yet it has not been explained with all its aspects and in full detail. As the question becomes experimentally relevant, we feel that it is important to display all the subtleties involved. The paper will be organized as follows: in Sect. II we discuss the theoretical evaluation of ΔM_K , ΔM_{B_d} , and ΔM_{B_s} together with $\Delta\Gamma_{B_s}$; in Sect. III we analyze the phase of $(q/p)\bar{\rho}$; in Sect. IV we address other conventions or proposals; in Sect. V we lay out a proper definition of the angles in the unitarity triangle before giving a summary in Sect. VI.

II. THE SIGN OF ΔM

For our discussion to be more transparent, we adopt a formalism and conventions which are applicable equally to the $K^0 - \bar{K}^0$ and the $B^0 - \bar{B}^0$ complexes.

Consider a neutral meson P carrying a quantum number $F = -1$; it can denote a K^0 or B^0 . The time evolution of a state being a mixture of P and $\bar{P}$ is given by

$$i\hbar \frac{\partial}{\partial t} \Psi(t) = \mathcal{H} \Psi(t) \quad (2.1)$$

where $\Psi(t)$ is restricted to the subspace of P and $\bar{P}$:

$$\Psi(t) = \begin{pmatrix} a(t) \\ \bar{b}(t) \end{pmatrix}. \quad (2.2)$$

Assuming **CPT** symmetry, the matrix $\mathcal{H}$ is given by

$$\mathcal{H} = \mathbf{M} - \frac{i}{2} \mathbf{\Gamma} = \begin{pmatrix} M_{11} - \frac{i}{2} \Gamma_{11} & M_{12} - \frac{i}{2} \Gamma_{12} \\ M_{12}^* - \frac{i}{2} \Gamma_{12}^* & M_{11} - \frac{i}{2} \Gamma_{11} \end{pmatrix}, \quad (2.3)$$

where

$$\begin{aligned} M_{11} &= M_P + \sum_n \mathcal{P} \left[\frac{|\langle n; out | H_{weak} | P \rangle|^2}{M_P - M_n} \right] \\ M_{12} &= \langle P | H_{SW} | \bar{P} \rangle + \sum_n \mathcal{P} \left[\frac{\langle P | H_{\Delta F=1} | n; out \rangle \langle n; out | H_{\Delta F=1} | \bar{P} \rangle}{M_P - M_n} \right] \\ \Gamma_{11} &= 2\pi \sum_n \delta(M_P - M_n) |\langle n; out | H_{\Delta F=1} | P \rangle|^2 \end{aligned}$$

$$\begin{aligned}
\Gamma_{22} &= 2\pi \sum_n \delta(M_P - M_n) |\langle n; out | H_{\Delta F=1} | \overline{P} \rangle|^2 \\
\Gamma_{12} &= 2\pi \sum_n \delta(M_P - M_n) \langle P | H_{\Delta F=1} | n; out \rangle \langle n; out | H_{\Delta F=1} | \overline{P} \rangle.
\end{aligned} \tag{2.4}$$

The coupled Schrödinger equations Eq.(2.1) are best solved by diagonalizing the matrix $\mathcal{H}$.

We find that

$$\begin{aligned}
|P_1\rangle &= p|P\rangle + q|\overline{P}\rangle, \\
|P_2\rangle &= p|P\rangle - q|\overline{P}\rangle,
\end{aligned} \tag{2.5}$$

are mass eigenstates with eigenvalues

$$\begin{aligned}
M_1 - \frac{i}{2}\Gamma_1 &= M_{11} - \frac{i}{2}\Gamma_{11} + \frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12}) \\
M_2 - \frac{i}{2}\Gamma_2 &= M_{11} - \frac{i}{2}\Gamma_{11} - \frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12})
\end{aligned} \tag{2.6}$$

as long as

$$\left(\frac{q}{p}\right)^2 = \frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}} \tag{2.7}$$

holds. Obviously, there are two solutions to this condition, namely

$$\frac{q}{p} = \pm \sqrt{\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}}}. \tag{2.8}$$

Choosing the negative rather than the positive sign in Eq.(2.8) is equivalent to interchanging the labels $1 \leftrightarrow 2$ of the mass eigenstates, see Eq.(2.5), Eq.(2.6).

This binary ambiguity is a special case of a more general one. For antiparticles are defined only up to a phase; adopting a different phase convention - *e.g.* going from $\mathbf{CP}|P\rangle = |\overline{P}\rangle$ to $\mathbf{CP}|P\rangle = e^{i\xi}|\overline{P}\rangle$ - will modify $M_{12} - \frac{i}{2}\Gamma_{12} \equiv \langle P | \mathcal{H} | \overline{P} \rangle$:

$$M_{12} - \frac{i}{2}\Gamma_{12} \rightarrow e^{i\xi} [M_{12} - \frac{i}{2}\Gamma_{12}] \tag{2.9}$$

and thus

$$\frac{q}{p} \rightarrow e^{-i\xi} \frac{q}{p}, \tag{2.10}$$

yet leave the combination $\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12})$ invariant. This is as it should be since the differences in mass and width

$$\begin{aligned} M_2 - M_1 &= -2\text{Re} \left(\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12}) \right) \\ \Gamma_2 - \Gamma_1 &= 4\text{Im} \left(\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12}) \right) \end{aligned} \quad (2.11)$$

being observables, have to be insensitive to the arbitrary phase of $\overline{P}$.

Some comments are in order to elucidate the situation:

- We can define the labels 1 and 2 such that

$$\Delta M \equiv M_2 - M_1 > 0 \quad (2.12)$$

is satisfied. Once this *convention* has been adopted, it becomes a sensible question whether

$$\Gamma_2 > \Gamma_1 \quad \text{or} \quad \Gamma_2 < \Gamma_1 \quad (2.13)$$

holds, *i.e.* whether the heavier state is shorter or longer lived.

- In the limit of **CP** invariance the two mass eigenstates are **CP** eigenstates as well, and we can raise another meaningful question: is the heavier state **CP** even or odd? Since **CP** invariance implies $\arg \frac{M_{12}}{\Gamma_{12}} = 0$, $\frac{q}{p}$ becomes a pure phase: $|\frac{q}{p}| = 1$. It is then convenient to adopt a phase convention s.t. M_{12} is real; it leads to $\frac{q}{p} = \pm 1$ and $\mathbf{CP}|P\rangle = \pm|\overline{P}\rangle$ as remaining choices.

– With $\frac{q}{p} = +1$ we have

$$\begin{aligned} |P_1\rangle &= \frac{1}{\sqrt{2}}(|P\rangle + |\overline{P}\rangle) \\ |P_2\rangle &= \frac{1}{\sqrt{2}}(|P\rangle - |\overline{P}\rangle) \end{aligned} \quad (2.14)$$

For **CP** $|P\rangle = |\overline{P}\rangle$, P_1 and P_2 are **CP** even and odd, respectively and therefore

$$\begin{aligned}
M_- - M_+ &= M_2 - M_1 = -2\text{Re}\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12}) \\
&= -2M_{12}
\end{aligned} \tag{2.15}$$

For $\mathbf{CP}|P\rangle = -|\overline{P}\rangle$, on the other hand, P_1 and P_2 switch roles; i.e. P_1 and P_2 are $\mathbf{CP}$ odd and even now. Thus

$$M_- - M_+ = M_2 - M_1 = 2M_{12} \tag{2.16}$$

– Alternatively we can set $\frac{q}{p} = -1$:

$$\begin{aligned}
|P_1\rangle &= \frac{1}{\sqrt{2}}(|P\rangle - |\overline{P}\rangle) \\
|P_2\rangle &= \frac{1}{\sqrt{2}}(|P\rangle + |\overline{P}\rangle)
\end{aligned} \tag{2.17}$$

while maintaining $\mathbf{CP}|P\rangle = |\overline{P}\rangle$; P_1 and P_2 are then $\mathbf{CP}$ odd and even, respectively. Accordingly

$$\begin{aligned}
M_- - M_+ &= M_1 - M_2 = 2\text{Re}\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12}) \\
&= -2M_{12}
\end{aligned} \tag{2.18}$$

– Eq.(2.15) and Eq.(2.18) on one side and Eq.(2.16) on the other do not coincide on the surface; yet, we will see below that the theoretical prediction for M_{12} changes sign depending on the choice of $\mathbf{CP}|P\rangle = \pm|\overline{P}\rangle$. Thus they all agree, of course.

- Within a given theory for the $P - \overline{P}$ complex, we can evaluate M_{12} , as discussed below. Yet some care has to be applied in interpreting such a result. For expressing mass eigenstates explicitly in terms of flavour eigenstates involves some conventions; see the examples above. Once we adopt a certain convention, we have to stick with it; yet our original choice cannot influence observables. It is instructive to trace how this comes about.

The *relative* phase between M_{12} and Γ_{12} on the other hand represents an observable quantity describing indirect $\mathbf{CP}$ violation. Therefore, we adopt the notation

$$M_{12} = \overline{M}_{12}e^{i\xi}, \quad \Gamma_{12} = \overline{\Gamma}_{12}e^{i\xi}e^{i\zeta}, \quad \text{and} \quad \frac{\Gamma_{12}}{M_{12}} = \frac{\overline{\Gamma}_{12}}{\overline{M}_{12}}e^{i\zeta} = re^{i\zeta} \quad (2.19)$$

The sign of $\overline{M}_{12}$ and $\overline{\Gamma}_{12}$ are fixed such that ξ , and $\xi + \zeta$ are restricted to lie between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$, respectively; *i.e.*, the real quantities $\overline{M}_{12}$ and $\overline{\Gamma}_{12}$ are a priori allowed to be *negative* as well as *positive*! A relative minus sign between M_{12} and Γ_{12} is of course physically significant, while the absolute sign is not. Yet, we will see that the absolute sign provides us with a useful bookkeeping device.

A. ΔM_K

The two kaon mass eigenstates can unambiguously be labelled by their lifetimes as K_L and K_S . One can then address the question how they differ in other properties: data reveal that (i) the K_L is ever so slightly heavier: $M_L > M_S$ and (ii) the K_S [K_L] is mainly **CP** even [odd]. We then adopt the conventions

$$\Delta M = M_2 - M_1, \quad \text{and} \quad \Delta \Gamma = \Gamma_1 - \Gamma_2 \quad (2.20)$$

which make both differences positive:

$$\Delta M_K = M_L - M_S > 0, \quad \Delta \Gamma_K = \Gamma_S - \Gamma_L > 0. \quad (2.21)$$

Since **CP** violation is very small in the K system - $\zeta_K = \arg(\Gamma_{12}^K/M_{12}^K) \ll 1$ - we deduce from Eq.(2.11) in the notation of Eq.(2.19):

$$\Delta M_K \simeq -2\overline{M}_{12}^K \quad \text{and} \quad \Delta \Gamma_K \simeq 2\overline{\Gamma}_{12}^K. \quad (2.22)$$

We will show below that the standard model box diagram indeed reproduces the ‘correct’, *i.e.* observed sign for ΔM_K (and – more surprisingly – even for $\Delta \Gamma_K$).

B. ΔM_{B_d}

The situation is qualitatively different here. While the two mass eigenstates will have different lifetimes, no useful experimental information exists on that. It is actually quite unlikely that such information will become available in the foreseeable future, since $\Delta \Gamma_{B_d}/\Gamma_{B_d}$

is estimated not to exceed the 1% level *. Thus we have to rely on a theoretical prediction of ΔM_{B_d} .

We can confidently predict for B_d mesons

$$|r_{B_d}| \ll 1. \quad (2.23)$$

Since we have to leading nontrivial order in r_{B_d}

$$\left. \frac{q}{p} \right|_{B_d} \simeq \sqrt{\frac{(M_{12}^{B_d})^*}{M_{12}^{B_d}}} \left(1 - \frac{r_{B_d}}{2} \sin \zeta_{B_d} \right), \quad (2.24)$$

we get from Eq.(2.11)

$$\Delta M_{B_d} = -2\overline{M}_{12}^{B_d} \quad \text{and} \quad \Delta \Gamma_{B_d} = 2\overline{\Gamma}_{12}^{B_d} \cos \zeta_{B_d}. \quad (2.25)$$

Note that the expression for ΔM_{B_d} is similar to the kaon case, albeit for a different reason, namely $|\Gamma_{12}/M_{12}| \ll 1$ rather than $\arg \frac{\Gamma_{12}}{M_{12}} \ll 1$; the latter quantity is actually not small for B_d mesons.

C. ΔM_{B_s} and $\Delta \Gamma_{B_s}$

There are two features that distinguish B_s from B_d decays in ways that are quite significant for our present discussion.

- While $r_{B_s} \ll 1$ holds also for B_s mesons, $\Delta \Gamma/\Gamma$ might not be that small. It has been estimated [8]

$$\frac{\Delta \Gamma}{\hat{\Gamma}}|_{B_s} \sim 0.18 \left(\frac{f_{B_s}}{200 \text{ MeV}} \right)^2, \quad \hat{\Gamma} = \frac{1}{2}(\Gamma_{B_{s,1}} + \Gamma_{B_{s,2}}). \quad (2.26)$$

Such a difference in the lifetimes of the two B_s mass eigenstates might actually become observable! This would raise the question whether the heavier state is longer or shorter lived.

*Remember that $\Gamma_S \gg \Gamma_L$ represents a kinematical accident!

- The $\Delta B = 2$ effective operator for B_s obtained from the box diagram is dominated by the quarks of the second and third families only. It thus predominantly conserves **CP** invariance making the two B_s mass eigenstates approximately **CP** eigenstates as well. This has two consequences:
 1. Rather than fit the decay rate evolution for $B_s \rightarrow D_s^{(*)}\pi$ or $B_s \rightarrow l\nu D_s^{(*)}$ with two separate exponentials, we can compare the lifetimes of the **CP** even eigenstate, as measured in $B_s \rightarrow \psi\eta$, with the average lifetime obtained from $B_s \rightarrow l\nu D_s^{(*)}$. We can also compare the lifetimes in $B_s \rightarrow \psi\phi$ for S- and P- wave final states being **CP** even and odd, respectively.
 2. We can raise the issue whether the **CP** even state is longer or shorter lived.

As for B_d mesons, we have

$$\left. \frac{q}{p} \right|_{B_s} = \sqrt{\frac{(M_{12}^{B_s})^*}{M_{12}^{B_s}}} \left(1 - \frac{r_{B_s}}{2} \sin \zeta_{B_s} \right) \quad (2.27)$$

and

$$\Delta M_{B_s} \simeq -2\overline{M}_{12}^{B_s} \quad \text{and} \quad \Delta \Gamma_{B_s} \simeq 2\overline{\Gamma}_{12}^{B_s} \quad (2.28)$$

where we have used the prediction that ζ_{B_s} , unlike ζ_{B_d} , is very small, since both M_{12} and Γ_{12} are dominated by contributions from the third and second family only.

D. Evaluating the sign of ΔM_K , ΔM_B , and $\Delta \Gamma_B$

After having established a notation equally convenient for the K and B (as well as D) case we calculate ΔM_K and ΔM_B . As is well known the dominant *short-distance* contribution to the effective $\Delta F = 2$ interaction within the Standard Model is obtained from the box diagram. One finds

$$\begin{aligned} \mathcal{H}_{eff}^{box}(\Delta F = 2, \mu) &= \left(\frac{G_F}{4\pi} \right)^2 M_W^2 \cdot \\ &\cdot \left[\eta_{cc}(\mu) \lambda_c^2 E(x_c) + \eta_{tt}(\mu) \lambda_t^2 E(x_t) + 2\eta_{ct}(\mu) \lambda_c \lambda_t E(x_c, x_t) \right] [\bar{q} \gamma_\mu (1 - \gamma_5) Q]^2 + h.c. \end{aligned} \quad (2.29)$$

where $Q = s, b$, and $q = d, s$; λ_i^Q denote combinations of KM parameters

$$\lambda_i^Q = \mathbf{V}_{iQ} \mathbf{V}_{iq}^*, \quad i = c, t, \quad (2.30)$$

and $E(x_i)$ and $E(x_i, x_j)$ reflect the box loops with equal and different internal quarks (charm or top), respectively:

$$E(x_i) = x_i \left(\frac{1}{4} + \frac{9}{4(1-x_i)} - \frac{3}{2(1-x_i)^2} \right) - \frac{3}{2} \left(\frac{x_i}{1-x_i} \right)^3 \log x_i \quad (2.31)$$

$$E(x_c, x_t) = x_c x_t \left[\left(\frac{1}{4} + \frac{3}{2} \frac{1}{1-x_t} - \frac{3}{4} \frac{1}{(1-x_t)^2} \right) \frac{\log x_t}{x_t - x_c} + (x_c \leftrightarrow x_t) - \frac{3}{4} \frac{1}{(1-x_c)(1-x_t)} \right]; \quad x_i = \frac{m_i^2}{M_W^2}. \quad (2.32)$$

The $\eta_{qq'}$ contain the QCD radiative corrections; they have been studied through next-to-leading level in order to understand the theoretical errors. We shall not go into the scale dependence as well as errors associated with uncertainties in Λ_{QCD} , m_t , etc. Such a discussion can be found in Ref. [10]. For us it is important to note that they are all *positive*.

These QCD corrections arise from evolving the effective Lagrangian from M_W down to the scale μ at which the hadronic expectation value is evaluated. The latter task is far from trivial even for a local four-fermion operator since on-shell matrix elements are controlled by long-distance dynamics. *In principle* the value of μ does not matter: the μ dependance of the $\Delta F = 2$ operator is compensated for by the μ dependance of the expectation value. *In practise* however, the available methods for calculating these matrix elements do not allow us to reliably track their μ dependance or they are applicable only for $\mu \sim 1$ GeV. Its size is customarily expressed as follows:

$$\langle P | (\bar{q} \gamma_\mu (1 - \gamma_5) Q) (\bar{q} \gamma_\mu (1 - \gamma_5) Q) | \bar{P} \rangle = -\frac{4}{3} B_P F_P^2 M_P, \quad \bar{P} = [Q \bar{q}] \quad (2.33)$$

which represents a parametrization rather than an ansatz as long as the value of B_P is left open.

$B_P = 1$ is referred to as *vacuum saturation* (VS) since it emerges when only the vacuum is inserted as intermediate state. Let us see the origin of the minus sign. One obtains

$$\langle P | (\bar{q}\gamma_\mu(1 - \gamma_5)Q)(\bar{q}\gamma_\mu(1 - \gamma_5)Q) | \bar{P} \rangle \Big|_{VS} = \frac{4}{3} \langle P | (\bar{q}\gamma_\mu(1 - \gamma_5)Q) | 0 \rangle \langle 0 | (\bar{q}\gamma_\mu(1 - \gamma_5)Q) | \bar{P} \rangle \quad (2.34)$$

upon inserting the vacuum intermediate state in the two types of box diagrams; this yields the colour factor $1 + 1/N_C = \frac{4}{3}$. Setting

$$\langle 0 | (\bar{q}\gamma_\mu(1 - \gamma_5)Q) | \bar{P}(k) \rangle = iF_P k_\mu \quad (2.35)$$

one finds with $\mathbf{CP} |0\rangle = |0\rangle$ and the convention $\mathbf{CP} |P\rangle = +|\bar{P}\rangle^\dagger$

$$\begin{aligned} \langle P | (\bar{q}\gamma_\mu(1 - \gamma_5)Q) | 0 \rangle &= \langle 0 | (\bar{Q}\gamma_\mu(1 - \gamma_5)q) | P \rangle^\dagger = \\ (\langle 0 | \mathbf{CP}^\dagger \mathbf{CP} (\bar{Q}\gamma_\mu(1 - \gamma_5)q) \mathbf{CP}^\dagger \mathbf{CP} | P \rangle)^\dagger &= -(\langle 0 | (\bar{q}\gamma_\mu(1 - \gamma_5)Q) | \bar{P} \rangle)^\dagger = iF_P k_\mu \end{aligned} \quad (2.36)$$

since

$$\mathbf{CP} \bar{Q}(\vec{x}) \gamma^\mu \gamma_5 q(\vec{x}) \mathbf{CP}^\dagger = -\bar{q}(-\vec{x}) \gamma^\mu \gamma_5 Q(-\vec{x}). \quad (2.37)$$

Several theoretical techniques have been employed to estimate the size of B_P . For a recent review, see Ref. [11]. Again for us it is important that they all yield $B_P > 0$.[‡] Within the Standard Model the short-distance contributions thus predict unequivocally

$$M_2 - M_1 > 0 \quad (2.38)$$

for K as well as B mesons. There are sizeable long distance contributions to ΔM_K , which are estimated to be positive as well [7]. In any case it would be quite contrived to expect them to cancel the short distance contributions.

[†]The VS result follows even with $\mathbf{CP} |0\rangle = e^{i\alpha} |0\rangle$.

[‡]The fudge factor B_P is sometimes called the bag factor since the MIT bag model was at an earlier time considered to yield a relatively reliable estimate for its size. It might be amusing to note while the bag model yields $B_B > 0$ as a very robust result, it is much less firm on B_K : both $B_K > 0$ as well as $B_K < 0$ emerge when varying the model parameters over a reasonable range!

With the definition $\mathbf{CP}|P\rangle = |\bar{P}\rangle$ used here, K_2 is the mainly $\mathbf{CP}$ odd state. Thus the Standard Model agrees with the experimental finding that K_L is heavier than K_S - a fact which is not often stressed in the literature.

For the K meson system, the box diagram does not lead to a reliable prediction for $\Delta\Gamma_K$ since the decay is dominated by $K \rightarrow \pi\pi$ which is anything but short distance dominated. The situation is quite different for $\Delta\Gamma_{B_s}$ which is driven by $[b\bar{s}] \rightarrow "c\bar{c}" \rightarrow [s\bar{b}]$ transitions.[§] It is reasonable to expect a short distance treatment to provide at least a semi-quantitative description. Hence we predict**

$$\Delta\Gamma_{B_s} = \Gamma_{B_s,even} - \Gamma_{B_s,odd} > 0. \quad (2.39)$$

III. THE PHASE OF $\frac{Q}{\bar{P}}\bar{\rho}(\psi K_S)$

Now that we know $\sin(\Delta Mt)$ describes the oscillations with positive ΔM , we need to know the phase of $\bar{\rho}(\psi K_S)$, which is defined by

$$\bar{\rho}(\psi K_S) = \frac{\langle \psi K_S | H_{\Delta B=1} | \bar{B} \rangle}{\langle \psi K_S | H_{\Delta B=-1} | B \rangle} = \frac{\langle \psi K_S | \psi \bar{K}^0 \rangle}{\langle \psi K_S | \psi K^0 \rangle} \frac{\langle \psi \bar{K}^0 | H_{\Delta B=1} | \bar{B} \rangle}{\langle \psi K^0 | H_{\Delta B=-1} | B \rangle}; \quad (3.1)$$

here we have used $\mathbf{CP}H_{\Delta B=1}\mathbf{CP}^\dagger = H_{\Delta B=-1}^*$.

Adopting the convention $\mathbf{CP}|P\rangle = +|\bar{P}\rangle$ we obtain

$$\begin{aligned} \langle \psi \bar{K}^0 | H_{\Delta B=1} | \bar{B}_d \rangle &= \langle \psi \bar{K}^0 | \mathbf{CP}^\dagger \mathbf{CP} H_{\Delta B=1} \mathbf{CP}^\dagger \mathbf{CP} | \bar{B}_d \rangle \\ &= -\langle \psi K^0 | H_{\Delta B=-1}^* | B_d \rangle \end{aligned} \quad (3.2)$$

where the minus sign reflects the fact that ψK^0 form a P wave. Thus

$$\bar{\rho}(\psi K_S) = -\frac{\langle \psi K_S | H_{\Delta B=-1}^* | B_d \rangle}{\langle \psi K_S | H_{\Delta B=-1} | B_d \rangle} = -\frac{\mathbf{V}_{cd}^* \mathbf{V}_{cs} \mathbf{V}_{cb} \mathbf{V}_{cs}^*}{\mathbf{V}_{cd} \mathbf{V}_{cs}^* \mathbf{V}_{cb}^* \mathbf{V}_{cs}} \quad (3.3)$$

[§]It is quite amusing that for $\Delta\Gamma_K$ the $s\bar{d} \rightarrow "u\bar{u}" \rightarrow d\bar{s}$ transition gives the correct sign for $\Delta\Gamma_K$.

**The question of the sign of $\Delta\Gamma_{B_d}$ might remain academic for experimental reasons. Theoretically the situation is not so clear due to large GIM cancellations [9].

where $V_{cd}^* V_{cs} / V_{cd} V_{cs}^* \sim \left(\frac{q_K}{p_K}\right)^*$. From Eq.(2.7), we see that

$$\frac{q}{p} = \sqrt{\frac{M_{12}^*}{M_{12}}} + \mathcal{O}(r) \propto \frac{\mathbf{V}_{tb}^* \mathbf{V}_{td}}{\mathbf{V}_{tb} \mathbf{V}_{td}^*} \quad (3.4)$$

Putting everything together we find the asymmetry is given by

$$\begin{aligned} \sin \Delta M_{Bt} \cdot \text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right) &= -\sin |\Delta M_{Bt}| \cdot \text{Im} \left(\frac{\mathbf{V}_{tb}^* \mathbf{V}_{td}}{\mathbf{V}_{tb} \mathbf{V}_{td}^*} \frac{\mathbf{V}_{cd}^* \mathbf{V}_{cb}}{\mathbf{V}_{cd} \mathbf{V}_{cb}^*} \right) \simeq \\ &\simeq \sin |\Delta M_{Bt}| \cdot \frac{2\eta(1-\rho)}{(1-\rho)^2 + \eta^2}, \end{aligned} \quad (3.5)$$

with the quantities η and ρ referring to the Wolfenstein representation of the CKM matrix.

With η inferred to be positive from ϵ one concludes that this asymmetry has to be *positive!*

IV. DIFFERENT CONVENTIONS

A. Using CP $|P\rangle = -|\bar{P}\rangle$

It is instructive to analyze what happens if one uses the equally allowed convention

$$\mathbf{CP} |P\rangle = -|\bar{P}\rangle. \quad (4.1)$$

Various intermediate expressions change; e.g., repeating the steps of Sect. II D one has

$$\langle P | (\bar{q} \gamma_\mu (1 - \gamma_5) Q) (\bar{d} \gamma_\mu (1 - \gamma_5) s) | \bar{P} \rangle \Big|_{V_S} = +\frac{4}{3} F_P^2 M_P, \bar{P} = [Q \bar{q}] \quad (4.2)$$

since now

$$\langle P | (\bar{q} \gamma_\mu (1 - \gamma_5) Q) | 0 \rangle = -i F_P k_\mu \quad (4.3)$$

That means that $\bar{M}_{12}$ changes sign; yet – as pointed out before – the two labels 1 and 2 exchange their roles then as well, see Eq.(2.16). The prediction that K_L is slightly heavier than K_S thus remains unaffected.

Furthermore the sign of $\bar{\rho}(\psi K_S)$ changes as well, see Sect. III. Thus the combination $\sin \Delta M_{Bt} \cdot \text{Im} \frac{q}{p} \bar{\rho}(\psi K_S)$ remains unchanged!

The ambiguity we have in the sign of ΔM_B is thus compensated by a corresponding ambiguity in the sign of $\frac{q}{p}\bar{\rho}(\psi K_S)$. We can see this also in the following way:

1. Changing $q/p \rightarrow -q/p$ maintains the defining property $(q/p)^2 = (M_{12}^* - \frac{i}{2}\Gamma_{12}^*)/(M_{12} - \frac{i}{2}\Gamma_{12})$, see Eq.(2.7), and cannot affect observables.
2. Yet the two mass eigenstates labeled by subscripts 1 and 2 exchange places, see Eq.(2.5).
3. The difference $\Delta M = -2\text{Re}[\frac{q}{p}(M_{12} - \frac{i}{2}\Gamma_{12})]$ then flips its sign - yet so does $\text{Im}\frac{q}{p}\bar{\rho}(\psi K_S)$!
4. The product $(\sin \Delta M_B t) \cdot \text{Im}\frac{q}{p}\bar{\rho}(\psi K_S)$ therefore remains invariant.

B. Another approach

The authors of [6] define

$$\Delta M_B = M_H - M_L \tag{4.4}$$

where H [L] stands for heavy [light]; ΔM is thus positive by definition. They also define

$$\begin{aligned} |B_L\rangle &= p|B^0\rangle + q|\bar{B}^0\rangle \\ |B_H\rangle &= p|B^0\rangle - q|\bar{B}^0\rangle \end{aligned} \tag{4.5}$$

For clarity, let us imagine a world in which **CP** is not violated. Then it is sensible to ask if the **CP** even state is heavier or lighter than the **CP** odd state. For good reason they have not decided on this question, since there is a freedom in the sign of

$$\frac{q}{p} = \pm \sqrt{\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}}} \tag{4.6}$$

If in computing the mass difference $M_2 - M_1$ turns out to be negative – yet we want to keep the convention **CP** $|P\rangle = |\bar{P}\rangle$ – we must choose the minus sign in Eq.(4.6). To say it differently: if we require $\Delta M_B > 0$ we have to compute the sign of $M_2 - M_1$ to decide on the sign for q/p in Eq.(4.6). Alternatively we can keep the plus sign if we adopt

CP $|B^0\rangle = -|\bar{B}^0\rangle$. In either case we must compute the sign of $M_2 - M_1$. The convention we must adopt is not fixed in this approach until we perform the theoretical computation of $M_2 - M_1$. We feel it is more natural to define a convention independent of prior theoretical computations.

FIGURES




FIG. 1. The unitarity triangle for B decays.

V. RELATION TO THE UNITARITY TRIANGLE

To conclude our discussion we explicitly restate the connections between the angles in the unitarity triangle, the CKM parameters and observable **CP** asymmetries.

The angles ϕ_1 , ϕ_2 and ϕ_3 are defined in Fig.1; as can be read off from Fig.2 they can be expressed by

$$\begin{aligned}\phi_1 &= \pi - \arg \left(\frac{-\mathbf{V}_{tb}^* \mathbf{V}_{td}}{-\mathbf{V}_{cb}^* \mathbf{V}_{cd}} \right), \\ \phi_2 &= \arg \left(\frac{\mathbf{V}_{tb}^* \mathbf{V}_{td}}{-\mathbf{V}_{ub}^* \mathbf{V}_{ud}} \right), \\ \phi_3 &= \arg \left(\frac{\mathbf{V}_{ub}^* \mathbf{V}_{ud}}{-\mathbf{V}_{cb}^* \mathbf{V}_{cd}} \right).\end{aligned}\tag{5.1}$$

The **CP** asymmetry in $B_d \rightarrow \psi K_S$, which is driven by a single $\Delta B = 1$ operator, is given by

$$\begin{aligned}A_{\psi K_S} &= \sin(\Delta M_B t) \text{Im} \left(\frac{q}{p} \bar{\rho}(\psi K_S) \right) = \\ &= \sin(2\phi_1) \sin(|\Delta M_B| t)\end{aligned}\tag{5.2}$$

To the degree we can eliminate the Penguin contribution to $B_d \rightarrow \pi^+ \pi^-$ by, say, extracting the asymmetry for the isopin-two $\pi\pi$ final state, we have likewise:

$$A_{(\pi\pi)_{I=2}} = \sin(\Delta M_B t) \text{Im} \left(\frac{q}{p} \bar{\rho}([\pi\pi]_{I=2}) \right) =$$

$$= \text{Im} \left(\frac{V_{tb} V_{td}^* V_{ud} V_{ub}^*}{V_{tb}^* V_{td} V_{ud}^* V_{ub}} \right) \sin(|\Delta M_B t|) \simeq \sin(|\Delta M_B t|) \sin(2\phi_2) \quad (5.3)$$




FIG. 2. The unitarity triangle.

VI. SUMMARY

We have shown that the *sign* of the **CP** asymmetry in $B_d \rightarrow \psi K_S$ can be predicted within a *given theory* for ΔM_B and $\frac{q}{p}\bar{\rho}(\psi K_S)$ even if the sign of ΔM_B cannot be determined experimentally. The same holds for $B_d \rightarrow \pi\pi$ etc. if one knows the weak operators driving these transitions.

En passant we have reminded the reader that the Standard Model can reproduce the observed sign of ΔM_K (and roughly its magnitude) through *short-distance* dynamics ^{††}.

Our discussion illustrates that we can choose any convention we want – provided care is applied in treating everything consistently. We view it, however, as very useful to have a formalism for describing B oscillations that parallels that for kaons. Furthermore it is much more natural to invoke theory to decide on the sign of ΔM_B .

Acknowledgements

Work of IIB has been supported in part by the NSF under the grant number PHY 96-0508. Work of AIS has been supported in part by Grant-in-Aid for Special Project Research (Physics of **CP** violation).

^{††}In the charm complex on the other hand ΔM_D is dominated by long distance dynamics within the Standard Model.

REFERENCES

- [1] A. B. Carter and A. I. Sanda, *Phys. Rev.* **D23** 1567 (1981) .
- [2] I. I. Bigi and A. I. Sanda, *Nucl. Phys.* **B193** 85 (1981)
- [3] K. Akerstaff, *et al.*, *Eur.Phys.J.***C5** 379,(1998).
- [4] CDF Collaboration (T.J. LeCompte for the collaboration). FERMILAB-CONF-98-293-E. To be published in the proceedings of Workshop on CP Violation, Adelaide, Australia, 3-8 Jul 1998.
- [5] Y. Grossman, B.Kayser and Y. Nir, *Phys.Lett.* **B 415** 90 (1997).
- [6] Y. Nir and H. R. Quinn, in *B decays* Edited by S. Stone World Scientific, Singapore, 1994.
- [7] I.I. Bigi and A.I. Sanda, *Phys. Lett.* **148B** (1984) 205-210.
- [8] M. Voloshin, M. Shifman, N. Uraltsev, and V. Khoze *Sov. J. Nucl. Phys.* **46** 112, (1987).
- [9] I.I. Bigi, V. Khoze, A. Sanda and N. Uraltsev, in: *CP Violation* Edited by C. Jarlskog, World Scientific, Singapore (1988).
- [10] G. Buchalla, A. J.Buras, and M. E. Lautenbacher, *Nucl. Phys.* **B370** 69 (1992) ; A. J. Buras, M. Jamin, and M. E. Lautenbacher, *Nucl. Phys.* **B408** 209 (1993) .
- [11] For a comprehensive review on hadronic matrix elements see: A. Soni, *Nucl. Phys.* **B47** (Proc. Suppl.)43 (1996) .